















- Sundarbans in Bangladesh and India using multi-temporal satellite data from 1973 to 2000. *Estuarine, Coastal and Shelf Science* 73: 91-100.
- [6] Datta D, Deb S. (2012). Analysis of coastal land use/land cover changes in the Indian Sunderbans using remotely sensed data. *Geo-spatial Information Science* 15: 241-250.
- [7] Giri C, Long J, Abbas S, Murali R, Qamer FM, Pengra B, Thau D. (2014). Distribution and dynamics of mangrove forests of South Asia. *Journal of Environmental Management* 148: 1-11.
- [8] Morawitz DF, Blewett TM, Cohen A. (2006). Using NDVI to assess vegetative land cover change in central puget sound. *Environ. Monit. Assess* 114(1-3): 85-106.
- [9] Nandy S, Kushwaha SPS. (2011). Study on the utility of IRS 1D LISS-III data and the classification techniques for mapping of Sunderban mangroves. *J. Coast. Conserv.* 15: 123-137.
- [10] Rahman MM, Ullah M, Lan M, Sumantyo JT, Kuze H, Tateishi R. (2013). Comparison of Landsat image classification methods for detecting mangrove forests in Sundarbans. *Int. J. Remote Sens.* 34: 1041-1056.
- [11] Raha AK, Zaman S, Sengupta K, Bhattacharyya SB, Raha S, Banerjee K, Mitra A. (2013). Climate change and sustainable livelihood programme: A case study from Indian Sundarbans. *J. Ecol.* 107: 335-348.
- [12] Raha A, Das S, Banerjee K, Mitra A. (2012). Climate change impacts on Indian Sunderbans: A time series analysis (1924-2008). *Biodivers. Conserv.* 21: 1289-1307.
- [13] Unnikrishnan AS, Kumar MRR, Sindhu B. (2011). Tropical cyclones in the Bay of Bengal and extreme sea-level projections along the east coast of India in a future climate scenario. *Curr. Sci. India* 101: 327-331.
- [14] Hazra S, Ghosh T, DasGupta R, Sen G. (2002). Sea level and associated changes in the Sundarbans. *Sci. C.* 68: 309-321.
- [15] Nandy S, Bandopadhyay S. (2011). Trend of sea level change in the Hugli estuary, India. *Indian J. Geo-Mar. Sci.* 40: 802-812.
- [16] Jayappa KS, Mitra D, Mishra AK. (2006). Coastal geomorphological and land-use and land cover study of Sagar Island, Bay of Bengal (India) using remotely sensed data. *Int. J. Remote Sens.* 27(17): 3671-3682.
- [17] Mitra D, Karmekar S. (2010). Mangrove classification in sunderban using high resolution multi spectral remote sensing data and GIS. *Asian J. Environ. Disast. Manage.* 2(2): 197-207.
- [18] Thomas JV, Arunachalam A, Jaiswal R, Diwakar PG, Kiran B. (2014). Dynamic land use and coastline changes in active estuarine regions – a study of sunderban delta, the international archives of the photogrammetry. *Remote Sensing and Spatial Information Sciences*, Volume XL-8, 2014 ISPRS Technical Commission VIII Symposium, Hyderabad, India.