

REFERENCES

- [1] Abaje IB, Ishaya S, Usman SU. (2010). An analysis of rainfall trends in Kafanchan, Kaduna State, Nigeria. *Research Journal of Environmental and Earth Sciences* 2(2): 89-96.
- [2] Abubakar SM, Kudanmiya YR, Eyong PN. (2002). Assessment of environmental degradation using satellite remote sensing technologies in Talata Mafara Area, Northern Nigeria. *Environmental Review* 4(1): 577-586.
- [3] Ajileye OO, Aigbiremolen IM, Mohammed SO, Halilu AS, Alaga AT. (2016). Effect of climatic variability on drought occurrence probability over Nigeria. *British Journal of Applied Science and Technology* 12(1): 1-13.
- [4] Ali IN. (2013). Fuel wood and vegetation change in Northern Nigeria: An exploration using remote sensing (RS), geographical information systems (GIS) and field reports. Unpublished PhD Thesis. University of Portsmouth, UK.
- [5] Ati OF, Tabitha Sheyin, Innocent Abbas, Seidu O, Mohammed. (2010). Assessing changes in Kagoro Forest, Kaduna State Nigeria using remote sensing and GIS. *Res. J. Appl. Sci. Eng. Technol.* 2(2): 121-132.
- [6] Audu EB. (2013). Fuel wood consumption and desertification in Nigeria. *International Journal of Science and Technology* 3(1): 1-5.
- [7] Baburao K, Ayse K, Kenneth H. (2013). Estimating crop coefficients using remote sensing-based vegetation index. *Remote Sens.* 5: 1588-1602. <http://doi.org/10.3390/rs5041588>
- [8] Zaharaddeen I, Mohammed SO, Auwal FA, Abel GR. (2017). Estimation of biophysical properties in lower river Kaduna catchment area Kaduna, Nigeria. *Asian Journal of Environment and Ecology (AJEE)* 4(1): 1-13. <http://doi.org/10.9734/AJEE/2017/32960>
- [9] Benkouider F, Abdellaoui A, Hamami L, Elaihar M. (2013). Spatio temporal analysis of vegetation by vegetation indices from multi-dates satellite images: Application to a semi-arid area in Algeria. *Energy Procedia* 36: 667-675.
- [10] El-Shirbeny MA, Aboelghar MA, Arafat SM, El-Gindy AG. (2014). Assessment of the mutual impact between climate and vegetation cover using NOAA-AVHRR and Landsat data in Egypt. *Arabian Journal of Geosciences* 7(4): 1287-1296.
- [11] El-Shirbeny MA, Ali AM, Badra MA, Bauomy EM. (2014). Assessment of wheat crop coefficient using remote sensing techniques. *World Research Journal of Agricultural Sciences* 1(2): 12-17.
- [12] Musa HD, Bukar S. (2010). Integrated remote sensing approach to desertification monitoring in the crop-rangeland area of Yobe State, Nigeria. *Journal of Sustainable Development in Africa* 12(5): 236-250.
- [13] Hatton T. (2001). Land use and catchment water balance. Technical Report 18/01. CSIRO Land and Water.
- [14] Hossein T, Ali A, Hossein ZPT, Shifteh SB. (2011). Spatial distribution and temporal variation of reference evapotranspiration in arid and semi-arid regions of Iran. *Hydrol. Process* 814: 1-13.
- [15] Justice CO, Townshend JRG. (2002). Special issue on the moderate resolution imaging spectroradiometer (MODIS): A new generation of land surface monitoring. *Remote Sens. Environ* 83: 1-2.
- [16] Kintada NR, Gurrum MK, Eadara A, Velagala VR. (2014). Land use/ land cover and NDVI analysis for monitoring the health of micro-watersheds of sarada river basin, Visakhapatnam District, India. *J Geol Geosci* 3: 146. <http://doi.org/10.4172/2329-6755.1000146>
- [17] Mohammed AA, Aliyu DG. (2014). Urban Vegetation Study of Kaduna Metropolis using GIS and remotely sensed Data. *Journal of Natural Sciences Research* 4(2): 160-171.
- [18] NASA. Landsat 7 Science Data Users Handbook (United States). https://landsat.gsfc.nasa.gov/wp-content/uploads/2016/08/Landsat7_Handbook.pdf
- [19] NASA. Landsat 8 Science Data Users Handbook (United States). https://landsat.usgs.gov/documents/Landsat8DataUsers_Handbook.pdf
- [20] Ndabula C, Averik PD, Jidauna GG, Abaje I, Oyatayo TK, Iguisi EO. (2013). Analysis of the Spatio-temporal dynamics of land use/ land cover structures in the kaduna inner core city region, Nigeria. *American Journal of Environmental Protection* 1(4): 112-119. <http://doi.org/10.12691/env-1-4-7>
- [21] Chris N. (2006). Assessment of land use/Landover changes in Kaduna Metropolitan area using remote sensing and geographic information system techniques. Unpublished Msc Thesis, Abu Zaria.
- [22] Ogutu JO, Piepho HP, Dublin HT, Bhola N, Reid RS. (2007). El Niño-southern oscillation, rainfall, temperature and normalized difference vegetation index fluctuations in the mara-serengeti ecosystem. *Afr. J. Ecol.* 46: 132-143.
- [23] Syeda RS, Amjed A, Ashfaq A, Muhammad M, Zia-Ul-Haq M, Shakeel A, Sezai E, Hawa ZEJ. (2014). Normalized difference vegetation index as a tool for wheat yield estimation: A case study from Faisalabad, Pakistan. *Scientific World Journal* 725326: 8. <http://dx.doi.org/10.1155/2014/725326>
- [24] Taiye OA, Emmanuel AO. (2015). Sustainability of fuel wood harvesting from Afaka forest reserve, Kaduna State, Nigeria. *Journal of Agricultural Science* 7(1): 129-135. <http://dx.doi.org/10.5539/jas.v7n1p129>
- [25] Zaharaddeen I. (2017). Assessment of Spatio-Temporal variation of Evapotranspiration in Middle River Kaduna Catchment area, Nigeria. Unpublished M.Sc. Thesis, Department of Geography Kaduna State University 1-113.
- [26] Huete AR. (1988). A soil-adjusted vegetation index (SAVI). *Remote Sensing of Environment* 25: 295-309.
- [27] Liu Y, Wang J. (2015). The research and application of landslide surface crack monitoring method based on laser ranging mode. *Environmental and Earth Sciences Research Journal* 2(2): 19-24 <http://dx.doi.org/10.18280/eesrj.020204>
- [28] Yang G, Song G, Cui W, Baiyinbaoligao, Chen W. (2015). Analysis and evaluation of land use and landscape pattern changes in Caohai watershed. *Environmental and Earth Sciences Research Journal* 2(2): 1-6. <http://dx.doi.org/10.18280/eesrj.020201>