











- interactions. In: Davies, R.A., Elington, R.L. (eds). Beach Nearshore sedimentation. Soc. Of Economic Paleontologists and Mineralogists Special Pub. 24, Tulsa.
- [17] Carling, P.A. (1982). Temporal and spatial variation in intertidal sedimentation rates. *Sedimentology*, 29: 17-23. <https://doi.org/10.1111/j.1365-3091.1982.tb01705.x>
- [18] Adefolalu, D.O. (1981). The weather and climate of Calabar: a study of coastal microclimatology in equatorial tropics. Senate Research Project Report, University of Calabar.
- [19] Richter, R. (1952). Fluidal-Textur in Sediment – Gesteinen und über Sedifluktion überhaupt. *Notizblatt des Hessischen Landesamtes für Bodenforschung zu Wiesbaden*, 3(6): 67-81.
- [20] Frey, R.W. (1975). The realm of ichnology, its strengths and limitations. In: *The Study of Trace fossils*, (Ed. R.W. Frey) Springer -Verlag, New York, pp. 13-38.
- [21] Bhattacharya, A., Sarkar, S.K., Bhattacharya, A. (2003). An assessment of coastal modification in the low – lying tropical coast of Northeast India and role of natural and artificial forcings. International Conference on Estuaries and Coasts November 9-11, 2003, Hangzhou, China.
- [22] Guza, R.T., Inman, D.L. (1975). Edge waves and beach cusps. *J. Geophys. Res.*, 80: 2997-3012.
- [23] Chakrabarti, P. (1990). Process-response system analysis in macrotidal and mesotidal coastal plain of Eastern India. *Memor Geol. Surv. India*, 2: 165-187.
- [24] Wright, L.D., Short, A.D. (1984). Morphodynamics variability of surf zones and beaches: A synthesis. *Marine Geology*, 32: 93-118.